

tors increases or decreases as $t \rightarrow \infty$, one may observe fractal clustering in region 1. The fractal dimension d_L is determined by the history of these factors. It can be computed in terms of ‘Lyapunov exponents’ [14, 16].

Figs. 1a,b show both mechanisms at work: at large values of St , (region 1 in Fig. 1c) there is no discernible influence of the instantaneous $\mathbf{u}(\mathbf{r}_t, t)$ on the particle distribution. At small St , (region 2 in Fig. 1c), by contrast, the particles are seen to avoid regions of high vorticity (Fig. 1b, similar to Fig. 8 in [17]). In short, in limiting cases (regions 1 and 2 of Fig. 1c) the mechanisms of clustering are understood. But how ergodic and non-ergodic effects compete in the major part of the phase diagram Fig. 1c is not known (in particular not for the experimentally most relevant region 3 where Ku and St are of order unity). In [18] non-ergodic effects were characterised by correlating the degree of clustering with the probability of particles avoiding rotational regions of the flow. The interpretation of these numerical results, however, is complicated by the fact that this probability is significantly enhanced even when clustering is weak. In order to understand the importance of non-ergodic and ergodic effects, an analytical theory is required, valid at finite Stokes and Kubo numbers.

Summary. Here we derive a perturbation expansion for the Lyapunov exponents of particles in random flows, valid at finite St and Ku . We compute the Lyapunov fractal dimension d_L , and characterise fractal clustering in terms of the ‘dimension deficit’ $\Delta_L = d - d_L$. For particles suspended in two-dimensional random Gaussian in-

compressible flows, the new theory yields reliable results for $Ku < 0.2$ for arbitrary values of St , region 4 in Fig. 1c (schematic). We find, first, that for small values of St , the ‘centrifuge’ mechanism dominates, and $\Delta_L = 6Ku^2St^2$, consistent with [20, 21]. Second, in the limit of $Ku \rightarrow 0$ at finite values of St , non-ergodic effects remain important. Third, in region 1 of Fig. 1c, clustering is found to be entirely due to ergodic ‘multiplicative amplification’ [14], and $\Delta_L = 12\epsilon^2 \propto St$. Fourth, in general we find that the ergodic mechanism contributes substantially to clustering, unless St is very small. Fifth we show by numerical simulations of the model that at $Ku \sim 1$ and $St \sim 1$, ergodic ‘multiplicative amplification’ makes a substantial contribution to the observed clustering.

Method and results. Eq. (1) cannot be explicitly solved, since \mathbf{u} depends upon the particle position at time t . The implicit solution of Eq. (1) becomes (dimensionless variables $\mathbf{r}' = \mathbf{r}/\eta$, $t' = t/\tau$, $\mathbf{v}' = \mathbf{v}/u_0$, and $\mathbf{u}' = \mathbf{u}/u_0$)

$$\delta\mathbf{r}_t \equiv \mathbf{r}_t - \mathbf{r}_0 = Ku \left[St(1 - e^{-t/St})\mathbf{v}_0 + St^{-1} \int_0^t dt_1 \int_0^{t_1} dt_2 e^{-(t_1-t_2)/St} \mathbf{u}(\mathbf{r}_{t_2}, t_2) \right]. \quad (2)$$

Here and in the following the primes are omitted. We seek an approximate solution by expanding $\mathbf{u}(\mathbf{r}_t, t)$ in powers of $\delta\mathbf{r}_t$. Since according to Eq. (2), $\delta\mathbf{r}_t$ is of order Ku , iteration generates an expansion of $\mathbf{u}(\mathbf{r}_t, t)$ in powers of Ku . To second order, for example, we find

$$\begin{aligned} u_\alpha(\mathbf{r}_t, t) = & u_\alpha(\mathbf{r}_0, t) + \frac{Ku}{St} \int_0^t dt_1 \int_0^{t_1} dt_2 e^{-(t_1-t_2)/St} \sum_\beta \frac{\partial u_\alpha}{\partial r_\beta}(\mathbf{r}_0, t) u_\beta(\mathbf{r}_0, t_2) \\ & + \frac{Ku^2}{St^2} \int_0^t dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 e^{-(t_1-t_2+t_3-t_4)/St} \sum_{\beta, \delta} \frac{\partial u_\alpha}{\partial r_\beta}(\mathbf{r}_0, t) \frac{\partial u_\beta}{\partial r_\delta}(\mathbf{r}_0, t_2) u_\delta(\mathbf{r}_0, t_4) \\ & + \frac{1}{2} \frac{Ku^2}{St^2} \int_0^t dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 e^{-(t_1+t_2-t_3-t_4)/St} \sum_{\beta, \delta} \frac{\partial^2 u_\alpha}{\partial r_\beta \partial r_\delta}(\mathbf{r}_0, t) u_\beta(\mathbf{r}_0, t_3) u_\delta(\mathbf{r}_0, t_4) + O(Ku^3). \end{aligned} \quad (3)$$

Here Greek indices denote the components of \mathbf{u} , and for our purposes \mathbf{v}_0 can be set to zero. The coefficients in (3) are expressed in terms of \mathbf{u} and its derivatives at the fixed position \mathbf{r}_0 , with known statistical properties. This procedure can in principle be extended to any order in Ku , but in practice it is limited by the number of nested integrals appearing in (3) for higher orders. A C-program was written to symbolically evaluate the integrals. One may expand other functionals of the particle trajectories, such as the strain matrix $\mathbb{A}(\mathbf{r}_t, t)$ with elements $A_{\alpha\beta} = \partial u_\alpha / \partial r_\beta$. Previous analytical results on the clustering of inertial particles [14, 19] rest on the

‘ergodic assumption’ that the distribution of the strain matrix $\mathbb{A}(\mathbf{r}_t, t)$ at the particle position \mathbf{r}_t can be approximated by its distribution at \mathbf{r}_0 . This is satisfied in region 1, but what are the corrections outside this region? For the two-dimensional incompressible ($\text{Tr}\mathbb{A} = 0$) random flow described in Fig. 1 we find:

$$\begin{aligned} \overline{\text{Tr}\mathbb{A}^2} \equiv & \left\langle \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \text{Tr}\mathbb{A}^2(\mathbf{r}_t, t) \right\rangle = \frac{6Ku^2St}{(1+St)^2(1+2St)} \\ & - \frac{2Ku^4St(4 + 52St + 293St^2 + 548St^3 + 297St^4)}{(1+St)^4(2+St)(1+2St)^2(1+3St)} \end{aligned} \quad (4)$$

to order Ku^4 . The average in (4) consists of a long-time average along the particle trajectory \mathbf{r}_t , and an average over initial conditions \mathbf{r}_0 (denoted by $\langle \dots \rangle$). At finite values of the Kubo number, $\overline{\text{Tr}}\mathbf{A}^2$ differs from its ergodic average (which vanishes in incompressible flows): the dynamics is not strictly ergodic. In compressible flows, we find $\overline{\text{Tr}}\mathbf{A} \neq 0$ (the ergodic average still vanishes). This is consistent with a result [22] for the average strain in the advective limit of a one-dimensional (compressible) model. It was shown in [22] that the average strain must be taken into account to obtain the known result [9] for the advective Lyapunov exponent in this model.

The question is now: how does non-ergodicity affect the spatial distribution of the particles? The latter is characterised by the Lyapunov exponents of the particle flow, obtained by linearising Eq. (1). The maximal exponent λ_1 is given by

$$\lambda_1 = Ku \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \mathbf{n}_1(\mathbf{r}_t, t) \cdot \mathbb{Z}(\mathbf{r}_t, t) \mathbf{n}_1(\mathbf{r}_t, t), \quad (5)$$

$$\dot{\mathbb{Z}} = \text{St}^{-1}(\mathbf{A} - \mathbb{Z}) - Ku \mathbb{Z}^2, \quad \dot{\mathbf{n}}_1 = Ku(\mathbf{n}_2 \cdot \mathbb{Z} \mathbf{n}_1) \mathbf{n}_2. \quad (6)$$

Here \mathbf{n}_1 (\mathbf{n}_2) is the unit vector in the $\delta \mathbf{r}$ ($\dot{\delta \mathbf{r}}$)-direction,

and \mathbb{Z} is the matrix with elements $Z_{\alpha\beta} = \partial v_\alpha / \partial r_\beta$. In region 1, Eqs. (5,6) were solved in [14, 19]. At finite values of Ku , we compute λ_1 by generalising the procedure that led to Eq. (3). Starting from the implicit solution (2) of (1), and the implicit solutions of (6):

$$\mathbb{Z}(\mathbf{r}_t, t) = e^{-t/\text{St}} \mathbb{Z}(\mathbf{r}_0, 0) + \int_0^t dt_1 e^{-\frac{t-t_1}{\text{St}}} [\mathbf{A}(\mathbf{r}_{t_1}, t_1) / \text{St} - Ku \mathbb{Z}(\mathbf{r}_{t_1}, t_1)^2], \quad (7)$$

$$\mathbf{n}_1(\mathbf{r}_t, t) = \mathbf{n}_1(\mathbf{r}_0, 0) + Ku \int_0^t dt_1 [\mathbf{n}_2(\mathbf{r}_{t_1}, t_1) \cdot \mathbb{Z}(\mathbf{r}_{t_1}, t_1) \times \mathbf{n}_1(\mathbf{r}_{t_1}, t_1)] \mathbf{n}_2(\mathbf{r}_{t_1}, t_1), \quad (8)$$

we expand $\mathbf{A}(\mathbf{r}_t, t)$, $\mathbb{Z}(\mathbf{r}_t, t)$, $\mathbf{n}_1(\mathbf{r}_t, t)$, and $\mathbf{n}_2(\mathbf{r}_t, t)$ in powers of $\delta \mathbf{r}_t$. Iterating and averaging along particle trajectories as well as over initial conditions yields an expansion of λ_1 in powers of Ku , with St -dependent coefficients. The sum $\lambda_1 + \lambda_2 = Ku \overline{\text{Tr}} \mathbb{Z}$ is computed in a similar fashion. For particles in a two-dimensional incompressible random Gaussian flow (c.f. Fig. 1) we find

$$\lambda_1 = Ku^2 - Ku^4 \frac{6 + 16\text{St} + 16\text{St}^2 + 15\text{St}^3 + 5\text{St}^4}{(1 + \text{St})^3} + Ku^6 \left[\frac{1692 + 16464\text{St} + 68987\text{St}^2 + 165269\text{St}^3 + 258832\text{St}^4}{6(1 + \text{St})^5(2 + \text{St})^2(1 + 2\text{St})^2} + \frac{301534\text{St}^5 + 296820\text{St}^6 + 247404\text{St}^7 + 153480\text{St}^8 + 62136\text{St}^9 + 14400\text{St}^{10} + 1440\text{St}^{11}}{6(1 + \text{St})^5(2 + \text{St})^2(1 + 2\text{St})^2} \right], \quad (9)$$

$$\lambda_1 + \lambda_2 = -6Ku^4 \frac{\text{St}^2(1 + 3\text{St} + \text{St}^2)}{(1 + \text{St})^3} + 2Ku^6 \text{St}^2 \frac{8 + 92\text{St} + 598\text{St}^2 + 2509\text{St}^3 + 5760\text{St}^4 + 7176\text{St}^5 + 5052\text{St}^6 + 2076\text{St}^7 + 480\text{St}^8 + 48\text{St}^9}{(1 + \text{St})^5(2 + \text{St})^2(1 + 2\text{St})^2}, \quad (10)$$

to order Ku^6 . This is our main result. Eq. (9) yields the ergodic expansion [19] $\lambda_1/(\gamma\tau) = 2\epsilon^2 - 20\epsilon^4 + 480\epsilon^6 + \dots$ in region 1.

As $\text{St} \rightarrow 0$, Eq. (10) reflects Maxey's non-ergodic centrifuge mechanism: According to (1), a particle is advected by an effective velocity field \mathbf{v} with compressibility $\nabla \cdot \mathbf{v} = \text{Tr} \mathbb{Z}$. Maxey's result [1, 2] is obtained by expanding $\mathbb{Z} \approx \mathbb{Z}^{(0)} + \mathbb{Z}^{(1)}\text{St}$ in Eq. (6). One finds: $\nabla \cdot \mathbf{v} = -Ku \text{St} \overline{\text{Tr}} \mathbf{A}^2|_{\text{St}=0}$ (note that $\mathbf{A}(\mathbf{r}_t, t)$ depends upon the Stokes number because the particle trajectory \mathbf{r}_t depends upon St). This result shows that particles tend to aggregate in regions of high strain or low vorticity. However, since the velocity field is homogeneous, this lowest-order term averages to zero in incompressible flows. Expanding \mathbb{Z} to second order in St , one finds $\overline{\nabla \cdot \mathbf{v}} = -Ku \text{St}^2 \partial_{\text{St}} \overline{\text{Tr}}(\mathbf{A}^2)|_{\text{St}=0}$. Inserting Eq. (4) yields

$\lambda_1 + \lambda_2 = Ku \overline{\nabla \cdot \mathbf{v}} = -6Ku^4 \text{St}^2 + 4Ku^6 \text{St}^2 + O(Ku^8)$, consistent with the $\text{St} \rightarrow 0$ limit of (10).

We conclude that our new expansion (9,10) of the Lyapunov exponents correctly describes the different clustering mechanisms in the advective and ergodic regions (Fig. 1c): the 'centrifuge' effect and ergodic 'multiplicative amplification'. More importantly, Eqs. (9,10) allow to determine how the relative importance of the two mechanisms depends on Ku and St , as follows.

In two-dimensional incompressible flows, the fractal dimension deficit is given by $\Delta_L = (\lambda_1 + \lambda_2)/\lambda_2$. Fig. 2a shows that the new theory explains the limiting cases in region 1 in the parameter plane ($\Delta_L = 6Ku^2\text{St} = 12\epsilon^2$), and in region 2 ($\Delta_L = 6Ku^2\text{St}^2$). The theory also explains the cross-over between these two behaviours and compares well with results of numerical simulations. In order to further increase the accuracy, more terms than




FIG. 2: **a** Fractal dimension deficit as a function of $\epsilon^2 = Ku^2St/2$. Numerical simulations of the model described in Fig. 1 ($Ku = 0.02$ (\diamond), $Ku = 0.05$ (\triangle), $Ku = 0.1$ (\circ), and $Ku = 0.2$ (∇); theory according to Eqs. (9,10), solid lines, and limiting behaviours $\Delta_L \propto St$ and $\Delta_L \propto St^2$ (dashed lines). **b** Same but for larger values of $\epsilon^2 = Ku^2St/2$. Numerics, $Ku = 0.1$ (\circ) and $Ku = 1$ (\square); theory, Eqs. (9,10), $Ku = 0.1$ (solid line). **c** Relative importance of non-ergodic and ergodic contributions (see text), symbols and parameters as in **b**.

computed in Eqs. (9,10) must be included. The series is likely to be asymptotic requiring re-summation, and there may be additional non-analytic contributions [14, 19]. We note that for a slightly different estimate of the fractal dimension (the ‘correlation dimension’), the lowest-order behaviour of the dimension deficit in region 2, $\Delta_C \propto St^2$, was computed using different methods in [9, 10, 23–25]. For the model described in Fig. 1, we find $\Delta_C = 12 Ku^2St^2$ to lowest order in Ku and in St in region 2, and $\Delta_C = 24 \epsilon^2$ in region 1 (see also [26]).

These results raise the question: how important are non-ergodic contributions to fractal clustering at larger values of St and at finite Kubo numbers? The answer is summarised in Fig. 2b,c showing the dimension deficit Δ_L compared to an ergodic approximation, Δ_L^{erg} . The latter incorporates finite-time correlations of the velocity field \mathbf{u} but neglects non-ergodic effects. Ergodic approximations to the Lyapunov exponents and to Δ_L (referred to as $\lambda_{1,2}^{\text{erg}}$ and Δ_L^{erg}) are obtained by expanding Eqs. (5,6) as before, but replacing $\mathbf{A}(\mathbf{r}_t, t)$ in (6) with $\mathbf{A}(\mathbf{r}_0, t)$. The resulting analytical expressions for $\lambda_{1,2}^{\text{erg}}$ are determined by the fluctuations of $\mathbf{A}(\mathbf{r}_0, t)$. This is in contrast to $\lambda_{1,2}$, Eqs. (9,10), which are determined by the fluctuations of $\mathbf{u}(\mathbf{r}_0, t)$ and its derivatives. The ergodic approximation allows for finite values of Ku and St but must fail in the limit $St \rightarrow 0$, since the ‘centrifuge’ mechanism is not accounted for. In particular, to lowest order in Ku the exponents λ_1^{erg} and λ_2^{erg} are found to depend upon St . The exact exponents (9,10) by contrast, are independent of St to lowest order in Ku : $\lambda_1 = Ku^2$ and $\lambda_2 = -Ku^2$. This implies in particular that earlier results for the Lyapunov exponents in region 1 [14, 19] are in fact exact to lowest order in Ku for arbitrary values of St . This is due to the cancellation of two errors: neglecting non-ergodic effects, and neglecting finite-time correlations.

Non-ergodic effects dominate the clustering when $(\Delta_L - \Delta_L^{\text{erg}})/\Delta_L$ is close to unity (they are negligible when this ratio is close to zero). We have determined $\lambda_{1,2}^{\text{erg}}$ and Δ_L^{erg} to order Ku^6 [27]. We have also performed

computer simulations of this ‘ergodic model’ by calculating monodromy matrices (Eq. (30) in [21]) with $\mathbf{A}(\mathbf{r}_0, t)$ evaluated at the fixed position ver_0 . Fig. 2c shows that for $Ku = 0.1$, non-ergodic effects dominate at small values of ϵ^2 but are negligible when clustering is largest, near the peak in Δ_L shown in Fig. 2b. Also shown are the analytical result for Δ_L derived from Eqs. (9,10), and the corresponding expansion of $(\Delta_L - \Delta_L^{\text{erg}})/\Delta_L$. We observe good agreement. For $Ku = 1$, by contrast, the first terms in the perturbation expansion (9,10) do not give reliable results. But computer simulations show that non-ergodic effects are present for the whole range of Stokes numbers displayed in Fig. 2c. However, Fig. 2c also clearly shows that both mechanisms contribute in region 3. We emphasise that ergodic clustering by ‘multiplicative amplification’ makes a substantial contribution in this region, $(\Delta_L - \Delta_L^{\text{erg}})/\Delta_L \approx 0.3$.

-
- [1] M. R. Maxey, *J. Fluid Mech.* **174** (1987) 441
 - [2] R. A. Shaw, *Annu. Rev. Fluid. Mech.* **35** (2003) 183.
 - [3] G. Falkovich, A. Fouxon and G. Stepanov, *Nature* **419** (2002) 151-154.
 - [4] M. Wilkinson, B. Mehlig and V. Bezuglyy, *Phys. Rev. Lett.* **97** (2006) 048501.
 - [5] A. Bracco *et al.*, *Phys. Fluids* **11** (1999) 2280
 - [6] M. Wilkinson, B. Mehlig and V. Uski, *Astrophys. J. Suppl. Ser.* **176** (2008) 484
 - [7] R. Reigada *et al.*, *Proc. R. Soc. Lond. B* **270** (2003) 875
 - [8] M. Pinsky and A. Khain, *Q. J. Met. Soc.* **123** (1997) 165
 - [9] E. Balkovsky, G. Falkovich, and A. Fouxon, *Phys. Rev. Lett.* **81** (2001) 2790
 - [10] J. Chun *et al.*, *J. Fluid Mech.* **536** (2006) 219
 - [11] E. Calzavarini *et al.*, *Phys. Fluids* **20** (2008) 040702
 - [12] M. Gibert, H. Xu, and E. Bodenschatz, arXiv:1002.3755
 - [13] M. Cencini *et al.*, *J. Turbulence* **7** (2006) 1
 - [14] K. P. Duncan *et al.*, *Phys. Rev. Lett.* **95** (2005) 240602.
 - [15] N. G. van Kampen, *Stochastic processes in physics and chemistry*, 2nd ed., North-Holland, Amsterdam 1981.
 - [16] J. Sommerer and E. Ott, *Science* **259** (1993) 351
 - [17] L. Chen, S. Goto, and J. C. Vassilicos, *J. Fluid Mech.* **553** (2006) 143
 - [18] J. Bec *et al.*, *Phys. Rev. Lett.* **98** (2007) 084502
 - [19] B. Mehlig and M. Wilkinson, *Phys. Rev. Lett.* **92** (2004) 250602
 - [20] J. Bec, *Phys. Fluids* **15** (2003) L81
 - [21] M. Wilkinson *et al.*, *Phys. Fluids* **19** (2007) 113303
 - [22] M. Wilkinson, arXiv:0911.2917
 - [23] L. I. Zaichik and V. M. Alipchenkov, *Phys. Fluids* **15** (2003) 1776
 - [24] R. H. A. Ijzermans, E. Meneguz, and M. W. Reeks, *J. Fluid Mech.* **653** (2010) 99
 - [25] G. Falkovich and Pumir, *Phys. Fluids* **16** (2004) L47
 - [26] M. Wilkinson, B. Mehlig, and K. Gustavsson, *Europhys. Lett.* **89** (2010), 50002
 - [27] K. Gustavsson and B. Mehlig, unpublished. We have obtained the ergodic series also in a different way: by generalising the one-dimensional approach described by M. Wilkinson, *J. Stat. Phys.* **139** (2010) 345.